

NOTE VI.

ON DERMAPTERA

COLLECTED IN JAVA BY MR. E. JACOBSON

BY

MALCOLM BURR, D. Sc. etc.

Mr. E. Jacobson, of Semarang, Java, has kindly sent to me for determination a small collection of Dermaptera from Java. I am able to recognize twenty-seven species; none of these are actually new, but several are interesting and the female of *Tomopygia abnormis* Borm. is described for the first time, this species being hitherto known only from de Bormans' unique type in the Brunner collection, now in the Hofmuseum, Vienna.

Order **DERMAPTERA.**Suborder **Forficulina.**Superfamily I. **PROTODERMAPTERA.**Family 1. **Pygidicranidae.**Subfamily 1. *Diplatyinae.*1. *Diplatys nigriceps* Kirby.

Goenoeng Merbaboe: July 1910, 1 ♂. — Wonosobo:
April 1909, 1 ♂ and 1 ♀.

Notes from the Leyden Museum, Vol. XXXIV.

2. *Diplatys* sp.

Wonosobo: April 1909, one larva with very long segmented caudal setae.

Subfamily 2. *Echinosomatinae*.3. *Echinosoma sumatranum* Haan.

Goenoeng Oengaran: Dec. 1909, 1 ♀ and 4 larvae.
A common Malayan species.

Family 2. Labiduridae.

Subfamily 1. *Psalinae*.4. *Anisolabis annulipes* Luc.

Telaga Mendjer: May 1909, 1 ♂. — Island of Krakatau:
May 1908, 1 ♂.

A cosmopolitan species.

5. *Anisolabis* sp.

Goenoeng Pangerango: Oct. 1908; Wonosobo: April 1909;
Semarang: Jan. 1910.

All these specimens are immature, and I refrain from attempting to name them.

6. *Gonolabis oblita* Burr.

Semarang: March 1910, ♂, ♀ and nymph. — Sron dol:
August 1909, ♀.

Subfamily 2. *Labidurinae*.7. *Labidura riparia* Pall.

Semarang: Dec. 1909, 1 ♀. — Djocja: Dec. 1909, 1 larva.
A cosmopolitan species.

8. *Nala lividipes* Dufour.

Semarang: Jan., Aug., Sept., Oct. and Nov., many specimens.
This species is common from the Mediterranean to the Malay Archipelago and to Cape Colony.

9. *Tomopygia abnormis* Borm. var.

Semarang: April 1910, 2 ♀♀.

Differs from type in having perfectly developed black elytra (and consequently concealed scutellum) and wings, which are long and dull brown in colour. The antennae, broken in the type, have 25 segments; 1 is rather short and thick, 2 minute; 3 cylindrical, slender, and quite as long as 1; 4, 5, 6 very short, cylindrical, no longer than broad, the rest gradually lengthening, almost cylindrical, but faintly subpyriform; dull brown.

Subfamily 3. *Platylabiinae*.10. *Platylabia major* Dohrn.

Goenoeng Oengaran: Oct. 1910, 1 ♀.

This is the species which I have always recorded as *Palex sparattoides* Borm. — de Bormans recorded it under that name and also as *Platylabia major* Dohrn. I now consider it to be the true *P. major* of Dohrn, who only described the female. The other species previously ranged in *Platylabia* are not congeneric with it, and fall into the extended genus *Chaetospania* Karsch.

Superfamily II. PARADERMAPTERA.

Family Apachyidae.

11. *Apachyus chartaceus* Haan.

Goenoeng Oengaran: Oct. 1910, 1 larva.

Superfamily III. EUDERMAPTERA.

Family 1. Labiidae.

Subfamily 1. *Spongiphorinae*.12. *Irdex nitidipennis* Borm.

Wonosobo: April 1909, 2 ♂♂ (macrolabious form). —
Goenoeng Oengaran: Sept., 2 ♀♀.

Superficially the forcipate form is extraordinarily like *Nala tenuicornis* Borm., but the simplest way to distinguish it promptly is by the presence, in the latter species, of a distinct, though not very strong, keel on the elytra.

13. *Marava grandis* Dubr.

Semarang: Nov. 1910, 1 ♂.

Subfamily 2. *Labiinae*.

14. *Labia mucronata* Stål.

Batavia: Jan., June and Dec. 1908, 4 ♂♂. — Goenoeng Oengaran: Sept. 1910, 2 ♀♀.

15. *Labia pilicornis* Motsch.

Semarang, Batavia and Wonosobo: Jan., April, May, June, July and November, 1 ♂, several ♀♀ and 1 larva.

16. *Labia curvicauda* Motsch.

Goenoeng Oengaran: September, ♂ and ♀. — Semarang: Jan. and June, 3 ♀♀. — Wonosobo: April, 1 ♀.
A variable cosmopolitan species.

17. *Prolabia nigrella* Dubr.

Goenoeng Oengaran: June 1910.

This specimen is of the colour-variety described by me under the name of *L. myrmeca* (Notes Leyden Mus. XXX, Note XIII, p. 96, 1908).

18. *Prolabia arachidis* Yers.

Semarang: Oct. and Dec. 1909, 1 ♂ and 1 ♀.

A cosmopolitan species.

19. *Chaetospania thoracica* Dohrn.

Goenoeng Oengaran: Sept., 1 ♀.

Family 2. Chelisochoidae.

Subfamily 1. *Chelisochinae*.20. *Chelisoches morio* Fabr.

Wonosobo in May, Batavia in Jan., June, Aug., Oct. and Dec., Goenoeng Oengaran in June, and Semarang in Febr., March, May, July, Aug., Sept., Oct., Nov. and Dec., many specimens.

This is an abundant and wide spread oriental species.

21. *Proreus simulans* Stål.

Semarang: January to November, and Batavia: October and December; several specimens.

A common oriental species.

22. *Enkrates elegans* Borm.

Wonosobo: April and May 1909, 5 ♂♂, 6 ♀♀ and 1 ♀ nymph.

These specimens all agree in coloration and structure with *E. elegans* of de Bormans, except in one particular, and that is the pygidium of the male. In *E. elegans* this organ is transverse, thick, with a gentle broad rounded emargination, the corner slightly produced to short blunt points. In one of these males the pygidium is typical, but in the other four it is different: it is broad, but the posterior margin is thickened, gently concave instead of convex, with a small, narrow median incision, and the angles are somewhat produced laterally, as the thickened margin protrudes beyond the sides.

In other respects, the smooth head, transverse pronotum, and the coloration, these males are indistinguishable, and they are associated with the females with linguiform pygidium, originally described by me as a distinct species under the name of *Apterygida lingua* (Termes. Füz. XXV, p. 486, pl. XX, fig. 8, 1902), and later definitely referred to *E. elegans* (vide Burr, Ann. Mag. N. H. (8) III, p. 256, 1909).

Notes from the Leyden Museum, Vol. XXXIV.

I am unable to separate the two forms in any respect beyond the form of the pygidium in the male, and the only way out of the difficulty, which I can see, is to admit dimorphism in the male pygidium.

Perhaps it is this creature which is described by de Bormans as *Ch. laetior* (Tierreich, Forf. p. 87, 1900), as his account of the female pygidium agrees with this species rather better than with the description of Dohrn, who only knew the female. Dohrn's type I have not yet seen.

23. *Hamaxas semiluteus* Borm.

Wonosobo: May 1909, 1 ♀.

Family 3. Forficulidae.

Subfamily 1. *Opisthocosmiinae*.

24. *Eparchus forcipatus* Haan.

Pangerango: Oct. 1908, 1 ♀.

25. *Eparchus* sp.

Wonosobo: April and May, 3 ♀♀.

This does not appear to be the female of *E. tenellus* and may very likely be new, but it is necessary to see the male.

26. *Eparchus tenellus* Haan.

Semarang: Dec. 1910, 1 ♀.

27. ?*Eparchus* sp. n.

Wonosobo: May 1910, 1 ♀.

Without the male, this species is indeterminable: it is very likely new.

Dover, October 1911.