

THE CONVOLVULACEAE OF MALESIA X¹⁾

Additional notes on the species from New Guinea with description of a new *Merremia*

S. J. VAN OOSTSTROOM

Rijksherbarium, Leiden

After the publication of my paper on the *Convolvulaceae* of New Guinea (Nova Guinea n.s. 6, 1955, 1—32) several new collections were made in this island. Some of these contain species up till then unknown from New Guinea, other give a better idea on the distribution within the island of species earlier found there. All specimens are in the Rijks-herbarium, Leiden.

Merremia emarginata (Burm. f.) Hallier f.; Ooststr., Blumea 3 (1939) 312; Fl. Mal. I, 4 (1953) 444; l.c. I, 5 (1958) 560.

West New Guinea. Merauke, sea level, Aug. 1954. Van Royen 4609.

Merremia aniseiifolia Ooststr., Blumea 12 (1964) 363.

West New Guinea. Kebar Valley, Andjai, alt. 600 m, Sept. 1959, Moll BW 9511 (type in L).

Merremia steenisii Ooststr., sp. nov. — Fig. 1.

Planta herbacea, dense cinereo-pilosa, caulis gracilibus, scandentibus. *Folia* breviter petiolata, petiolis 3—5(—7) mm longis, lineari-oblonga vel linearia, 5—7 cm longa, 5—8(—12) mm lata, basi rotundata, apice acuta, mucronulata, nervis lateralibus utrinque 6—8 ascendentibus, subtus prominentibus. *Inflorescentiae* axillares, pedunculatae, 1-florae; pedunculis c. 12—15 mm longis, gracilibus, pedicellis c. 10—15 mm longis; bracteis subulatis c. 5 mm longis. Sepala exteriora 2 lanceolato-ovata vel lanceolata, acuta, c. 12—13 mm longa, interiora 3 lanceolata, acuminata, c. 14 mm longa. Corolla infundibuliformis, 3.5 cm longa, fasciis mesopetalibus extus sparse pilosis. Stamina inclusa, filamentis c. 5—6 mm longis, c. 2 mm supra basin corollae insertis, basi papillosis; antheris maturis non vel vix contortis, c. 3 mm longis. Discus annularis. Ovarium pilosum; stylo inclusu, c. 8 mm longo.

Territory of New Guinea. Sepik Dist.: Wewak-Angoram area, road from Yangoru to Kworo, Middle Sepik plains, in grassland, only a single specimen, fl. Sept. 1959, Robbins 2316 (type in CANB).

¹⁾ Part I in Blumea 3 (1938) 62—94; II in 3 (1939) 267—371; III in 3 (1940) 481—582; IV in 5 (1943) 339—411; V in 5 (1945) 689—691; VI in 6 (1950) 337—348; VII in 7 (1952) 171—178; VIII (by R. D. Hoogland) in 7 (1952) 179—192, and IX (by R. D. Hoogland) in 7 (1953) 310—319.




Fig. 1. *Merremia steenisii*. — a: flowering stem; b—e: sepals 1—4, outside; f₁: sepal 5, outside; f₂: sepal 5, inside; g: pistil; h: stamen. — *M. aniseifolia*. — i: stamen. — (a: $\times \frac{1}{2}$; b—g: $\times 3$; h, i: $\times 4\frac{1}{2}$).

This new species is doubtless closely related to *Merremia aniseifolia* which I described a number of years ago from West New Guinea (Blumea 12, 1964, 363). As will be evident by comparison of the descriptions of both, it has many characters in common with that species, and it is quite possible that more material becoming available will throw some other light upon their mutual relations. However, for the time being I feel justified to keep them separate.

M. steenisii can be distinguished from *M. aniseifolia* at first sight by the totally different appearance of the plant, mainly caused by the dense hairiness of stems, leaves, peduncles, pedicels, and sepals. Other points of difference with *M. aniseifolia* may be found in the shape and texture of the leaves which are relatively narrower and thicker, in the more prominent nerves on the lower leaf surface, the absence of warts on the sepals, of which the outer ones are a little shorter than the inner, the slightly pilose instead of glabrous midpetaline bands of the corolla, the somewhat shorter stamens inserted nearer to the corolla base, and the not or scarcely twisted ripe anthers, which are strongly spirally twisted in *M. aniseifolia*. A difference is also found in the base of the filaments; in *M. aniseifolia* this bears a long row of short hairs on both sides, the two rows being parallel and connected arcuately above and decurrent on the corolla base beneath; in the new species these hair rows are much shorter (fig. 1, h and i).

I name this species after Prof. Dr. C. G. G. J. van Steenis, the retiring director of the Rijksherbarium, Leiden, who during many years has done so much for the advancement of the study of the Malesian flora.

***Merremia quinquefolia* (L.) Hallier f.; Ooststr., Blumea 3 (1939) 324; Fl. Mal. I, 4 (1953) 446.**

Territory of New Guinea. Morobe Dist.: Wau, alt. 400 ft, March 1965, Millar NGF 23866; Crooked Creek, Bulolo, alt. 2200 ft, July 1969, Streimann & Kairo NGF 44040.

Territory of Papua. Central Dist.: 3 m. W. of Mt. Lawes, alt. 100 ft, July 1962, Schodde 2628; Rubulogo Creek, 18 m. N. of Port Moresby, alt. 100 ft, April 1967, Pullen 6642; id., Gordon, Oct. 1969, Pulsford UPNG 175.

New to New Guinea.

***Merremia quinata* (R.Br.) Ooststr.; Ooststr., Fl. Mal. I, 4 (1953) 447; Nova Guinea n.s. 6 (1955) 17. — *Ipomoea quinata* R. Br.; Ooststr., Blumea 3 (1940) 581.**

Territory of Papua. Milne Bay Dist.: Cape Vogel Peninsula, Menapi, alt. 100 m, April 1953, Brass 22012; ibidem, N. coast, near Medino village, alt. 25 m, Sept. 1954, Hoogland 4697; Baniari Subdist., Aribi near Dogura, sea level, June 1965, Galore NGF 17564.

Known from the Territory of New Guinea.

***Merremia dissecta* (Jacq.) Hallier f.; Ooststr., Blumea 3 (1939) 328; Fl. Mal. I, 4 (1953) 448.**

Territory of Papua. Central Dist.: Kairuku Subdist., Rogers airstrip, 8 m. W. of Kanosia plantation, alt. 30 ft, July 1962, Derbyshire 651; Fairfax Harbour, sea level, Febr. 1965, Gillison NGF 22228.

New to New Guinea.

***Merremia pacifica* Ooststr.; Ooststr., Fl. Mal. I, 5 (1958) 560.**

Territory of Papua. Louisiade Archipelago, Rossel I., Jinju, alt. 80 m, Oct. 1956, Brass 28540.

Ipomoea pes-tigridis L.; Ooststr., Blumea 3 (1940) 504; Fl. Mal. I, 4 (1953) 467; Nova Guinea n.s. 6 (1955) 24.

West New Guinea. Merauke, sea level, Febr. 1956, Van der Sijde BW 4009.

New to New Guinea.

Ipomoea triloba L.; Ooststr., Blumea 3 (1940) 509; Fl. Mal. I, 4 (1953) 468; Nova Guinea n.s. 6 (1955) 24.

West New Guinea. Merauke, sea level, Febr. 1956, Van der Sijde BW 4019.

Territory of New Guinea. Morobe Dist.: Lac, alt. 30 ft, Sept. 1964, Henry NGF 20390; ibidem, Huon road, alt. 50 ft, Aug. 1962, Van Royen NGF 15063.

Known from the Territory of Papua.

Ipomoea tiliacea (Willd.) Choisy; Ooststr., Blumea 5 (1942) 233; Fl. Mal. I, 4 (1953) 469; Nova Guinea n.s. 6 (1953) 25; Fl. Mal. I, 5 (1958) 561.

Territory of Papua. Northern Dist.: Tufi Subdist., near Wanigela, alt. 5 m, Sept. 1954, Hoogland 4774.

Known from West New Guinea.

Ipomoea obscura (L.) Ker-Gawl.; Ooststr., Blumea 3 (1940) 519; Fl. Mal. I, 4 (1953) 471; Nova Guinea n.s. 6 (1953) 26.

West New Guinea. Hollandia, alt. 30 m, Nov. 1956, Van der Sijde BW 4113; ibidem, April 1959, De Wilde & Vervoort 405; ibidem, road to Lake Sentani, alt. 25 m, July 1954, Van Royen 4507.

Known from the Territory of Papua.

Ipomoea fimbriosepala Choisy; Ooststr., Fl. Mal. I, 5 (1958) 561.

Territory of New Guinea. E. Highl. Dist.: Goroka Subdist., near Gitunu village, between Bena Bena and Dumantina R., alt. c. 1500 m, June 1956, Hoogland & Pullen 5288. — Morobe Dist.: Wau Subdist., Bulolo Forestry School, in sec. grasslands, alt. c. 2500 ft, Sept. 1967, Kairo & Streimann NGF 30897.

Ipomoea cairica (L.) Sweet; Ooststr., Blumea 3 (1940) 542; Fl. Mal. I, 4 (1953) 478; Nova Guinea n.s. 6 (1955) 27.

West New Guinea. Hollandia, Kota Baru, alt. 100 m, Oct. 1954, Lam 7760; Manokwari, alt. 5 m, Nov. 1960, Moll BW 9793.

Known from the Territory of New Guinea.

Ipomoea alba L.; Ooststr., Blumea 3 (1940) 547; Fl. Mal. I, 4 (1953) 480; Nova Guinea n.s. 6 (1955) 28.

Territory of Papua. Milne Bay Dist.: Kwagira R., Peria Creek, alt. 50 m, Sept. 1953, Brass 24291.

Known from the Territory of New Guinea.

Ipomoea hederifolia L.; Ooststr., Nova Guinea n.s. 6 (1955) 28; Fl. Mal. I., 5 (1958) 563 — *I. angulata* Lamk.; Ooststr., Blumea 3 (1940) 553; Fl. Mal. I., 4 (1953) 481.

West New Guinea. Hollandia, alt. 3 m, March 1957, Kalkman BW 3675; ibidem, Ifar, alt. 450 m, Nov. 1956, Van den Assem 11; Amban near Manokwari, alt. 100 m, March 1959, Versteegh & Vink BW 8395.

Territory of New Guinea. Morobe Dist.: Bulolo Valley, alt. 3500 ft, April 1959, Womersley NGF 11019 — E. Highl. Dist.: Waghi Valley, Kondiu, July 1960, Borgmann 113.

Territory of Papua. Milne Bay Dist.: Cape Vogel Peninsula, Menapi, alt. 5 m, April 1953, Brass 22007; D'Entrecasteaux Is., Fergusson I., Deidei, Gomwa Bay, alt. 20 m, July 1956, Brass 27322.

New to New Guinea.