



# A phylogenetic study of *Dothiorella* and *Spencermartinsia* species associated with woody plants in Iran, New Zealand, Portugal and Spain

J. Abdollahzadeh<sup>1</sup>, A. Javadi<sup>2</sup>, R. Zare<sup>2</sup>, A.J.L. Phillips<sup>3</sup>

## Key words

*Botryosphaeriaceae*  
*Dothiorella*  
ITS  
phylogeny  
*Spencermartinsia*  
systematics  
taxonomy

**Abstract** *Dothiorella* and *Spencermartinsia* are two botryosphaeriaceous genera with dark 2-celled conidia and found in parasitic, saprophytic or endophytic association with various woody host plants. Based on ITS and EF1- $\alpha$  sequence data and morphology, eight new species are described from Iran, New Zealand, Portugal and Spain. Of these, five species are placed in *Dothiorella*, namely *D. iranica*, *D. parva*, *D. prunicola*, *D. sempervirentis* and *D. striata*, and three species belong to *Spencermartinsia* named as *S. citricola*, *S. mangiferae* and *S. plurivora*. An identification key to the species of each genus is provided.

**Article info** Received: 24 April 2013; Accepted: 1 November 2013; Published: 14 January 2014.

## INTRODUCTION

*Dothiorella* was established by Saccardo in 1880 (Crous & Palm 1999) and according to Sutton (1980) *Dothiorella pyrenophora* Sacc. is the type species. Over time, the generic concept of *Dothiorella* has been debated and interpreted in various different ways. The taxonomic history of *Dothiorella* has been explained in details by Sutton (1977), Crous & Palm (1999), and Phillips et al. (2008, 2013). Crous & Palm (1999) studied the holotype of *D. pyrenophora* and concluded that *Dothiorella* is a synonym of *Diplodia*. This concept was followed by Denman et al. (2000), Zhou & Stanosz (2001) and Slippers et al. (2004), who recognised only two groups in *Botryosphaeria* corresponding to the asexual genera, *Fusicoccum* and *Diplodia* with hyaline and pigmented mature conidia, respectively. However, based on morphology and molecular data, Phillips et al. (2005) resurrected *Dothiorella* for species with conidia that become brown and 1-septate while they are still attached to the conidiogenous cells, and reserved *Diplodia* for species with conidia that are hyaline and become dark and septate only some time after discharge from the pycnidia. They also provided emended descriptions for *Diplodia* and *Dothiorella*. Since, there is no culture or DNA sequence for the type specimen and no authentic isolate or culture is available for the type species of *Dothiorella*, it is necessary to designate an epitype and ex-epitype culture for this important species.

Phillips et al. (2008) in a polyphasic approach using morphology and multi-gene sequence data (SSU, LSU, ITS, EF1- $\alpha$  and  $\beta$ -tubulin) introduced *Spencermartinsia* for species having brown, 1-septate ascospores with an apiculus at either end and transferred *Dothiorella viticola* to *Spencermartinsia*. Species of *Dothiorella* also have brown, 1-septate ascospores, but can be

distinguished from *Spencermartinsia* by the absence of apiculi. In subsequent multi-gene phylogenetic analyses (Liu et al. 2012, Phillips et al. 2013) it was confirmed that *Dothiorella* and *Spencermartinsia* are two distinct genera in the *Botryosphaeriaceae*, though obviously closely related (see Slippers et al. 2013).

Species of *Dothiorella* and *Spencermartinsia*, as members of *Botryosphaeriaceae*, are known saprophytes, pathogens and endophytes in association with various woody plants. Until recently, like other members of *Botryosphaeriaceae*, species in *Dothiorella* were described based on host association, which led to the introduction of many species names. A search of Index Fungorum (March 2013; www.indexfungorum.org) lists 363 names in *Dothiorella*, while MycoBank lists 384 species names (accessed March 2013). Since, host association is not considered to be an important factor in species definition of the *Botryosphaeriaceae* (Slippers et al. 2004, 2013) most of these names are likely synonyms or they need to be transferred to *Spencermartinsia*.

A number of isolates morphologically resembling *Dothiorella* or *Spencermartinsia* were collected on different woody hosts in Iran, New Zealand, Portugal and Spain. The aim of this study was to characterise these isolates based on molecular data combined with morphology.

## MATERIALS AND METHODS

### Fungal isolates

Isolations were made by transferring conidia produced in pycnidia on twigs with canker or dieback symptoms to potato dextrose agar (1/2 PDA, Difco laboratories) or water agar (WA) supplemented with chloramphenicol (100 mg/L). After incubating overnight at 25 °C, single germinating conidia were transferred to fresh PDA plates and single conidium cultures were prepared for all isolates (Table 1).

### Morphology and culture characteristics

Sporulation was induced by growing isolates on 2 % WA supplemented with pieces of double-autoclaved halved poplar

<sup>1</sup> Department of Plant Protection, Faculty of Agriculture, University of Kurdistan, P.O. Box 416, Sanandaj, Iran;  
corresponding author e-mail: J.abdollahzadeh@uok.ac.ir.

<sup>2</sup> Department of Botany, Iranian Research Institute of Plant Protection, P.O. Box 1454, Tehran 19395, Iran.

<sup>3</sup> Centro de Recursos Microbiológicos, Departamento de Ciências da Vida, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal.

**Table 1** Isolates considered in this study.

Species	Culture no. <sup>1</sup>	Host	Location	Collector	GenBank numbers <sup>2,3</sup>	
					ITS	EF1- $\alpha$
<i>Dothiorella iranica</i>	<b>IRAN1587C</b> /CBS124722	<i>Olea europaea</i>	Iran, Golestan	A. Javadi	<i>KC898231</i>	<i>KC898214</i>
<i>D. parva</i>	<b>IRAN1579C</b> /CBS124720	<i>Corylus avellana</i>	Iran, Ardabil	J. Abdollahzadeh/A. Javadi	<i>KC898234</i>	<i>KC898217</i>
	IRAN1585C/CBS124721	<i>C. avellana</i>	Iran, Ardabil	J. Abdollahzadeh/A. Javadi	<i>KC898235</i>	<i>KC898218</i>
<i>D. pretoriensis</i>	<b>CMW36480</b>	<i>Acacia karroo</i>	South Africa	F. Jami/M. Gryzenhout	JQ239405	JQ239392
	CMW36481	<i>A. karroo</i>	South Africa	F. Jami/M. Gryzenhout	JQ239406	JQ239393
<i>D. prunicola</i>	CBS124723/ <b>IRAN1541</b>	<i>Prunus dulcis</i>	Portugal, Algarve	E. Diogo	EU673313	EU673280
<i>D. sempervirens</i>	IRAN1581C/CBS124719	<i>Cupressus sempervirens</i>	Iran, Golestan	M.A. Aghajani	<i>KC898237</i>	<i>KC898220</i>
	<b>IRAN1583C</b> /CBS124718	<i>C. sempervirens</i>	Iran, Golestan	M.A. Aghajani	<i>KC898236</i>	<i>KC898219</i>
	IRAN1580C	<i>C. sempervirens</i>	Iran, Golestan	M.A. Aghajani	n.s.	n.s.
	IRAN1582C	<i>C. sempervirens</i>	Iran, Golestan	M.A. Aghajani	n.s.	n.s.
	IRAN1586C	<i>C. sempervirens</i>	Iran, Golestan	M.A. Aghajani	n.s.	n.s.
<i>D. striata</i>	ICMP16819/CBS124730/ IRAN1503C	<i>Citrus sinensis</i>	New Zealand	S.R. Pennycook/P.R. Johnston	EU673320	EU673287
	<b>ICMP16824</b> /CBS124731/ IRAN1572C	<i>C. sinensis</i>	New Zealand	S.R. Pennycook/P.R. Johnston	EU673321	EU673288
<i>D. uruguayensis</i>	<b>UY672</b> /CBS124908	<i>Hexalaminis edulis</i>	Uruguay	C. Perez	EU080923	EU863180
<i>Dothiorella</i> sp.	IRAN1570C/CBS124717	<i>Juglans regia</i>	Iran, Kermanshah	J. Abdollahzadeh/A. Javadi	<i>KC898233</i>	<i>KC898216</i>
	IRAN1573C/CBS124716	<i>J. regia</i>	Iran, Jolfa	J. Abdollahzadeh/A. Javadi	<i>KC898232</i>	<i>KC898215</i>
	IRAN1576C	<i>J. regia</i>	Iran, Kermanshah	J. Abdollahzadeh/A. Javadi	n.s.	n.s.
	IRAN1577C	<i>J. regia</i>	Iran, Kermanshah	J. Abdollahzadeh/A. Javadi	n.s.	n.s.
<i>Spencermartinsia citricola</i>	ICMP16827/CBS124728/ IRAN1504C	<i>C. sinensis</i>	New Zealand	S.R. Pennycook/P.R. Johnston	EU673322	EU673289
	<b>ICMP16828</b> /CBS124729/ IRAN1505C	<i>C. sinensis</i>	New Zealand	S.R. Pennycook/P.R. Johnston	EU673323	EU673290
<i>S. mangiferae</i>	IRAN1545C	<i>Mangifera indica</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898223</i>	<i>KC898206</i>
	IRAN1546C/CBS124726	<i>M. indica</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898222</i>	<i>KC898205</i>
	<b>IRAN1584C</b> /CBS124727	<i>M. indica</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898221</i>	<i>KC898204</i>
<i>S. plurivora</i>	IRAN1537C	<i>Citrus</i> sp.	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898226</i>	<i>KC898209</i>
	IRAN1538C	<i>C. sempervirens</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898229</i>	<i>KC898212</i>
	IRAN1552C	<i>Casuarina equisetifolia</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898228</i>	<i>KC898211</i>
	IRAN1553C	<i>Malus domestica</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898227</i>	<i>KC898210</i>
	IRAN1556C/CBS124725	<i>Prunus armeniaca</i>	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898230</i>	<i>KC898213</i>
	<b>IRAN1557C</b> /CBS124724 CJA257	<i>Citrus</i> sp. <i>Eucalyptus</i> sp.	Iran, Hormozgan	J. Abdollahzadeh/A. Javadi	<i>KC898225</i> <i>KC898224</i>	<i>KC898208</i> <i>KC898207</i>

<sup>1</sup> Ex-type cultures are in bold face.

<sup>2</sup> Sequence numbers in italics were obtained in the present study. All others were retrieved from GenBank.

<sup>3</sup> n.s.: Not sequenced.

twigs. The plates were incubated at 25 °C under mixed black (nUV) and fluorescent light for 2–6 wk. Conidiomata were dissected and mounted in 100 % lactic acid. Observations and digital images were made with a light microscope and digital camera (Leica or Olympus). From measurements of 50 conidia the mean, standard deviation and 95 % confidence intervals were calculated. Dimensions are given as the range of measurements with extremes in brackets followed by 95 % confidence limits and mean  $\pm$  standard deviation. Dimensions of other structures are given as the range of at least 20 measurements. Colony morphology, colour (Rayner 1970) and growth rates between 5 and 35 °C in 5 °C intervals, were determined on 2 % malt extract agar (MEA, Difco laboratories) incubated in the dark.

#### DNA extraction, PCR amplification and sequencing

Genomic DNA was extracted from 4–7 d old cultures grown in 2 % malt extract broth (MEB) incubated at room temperature using the method as described by Abdollahzadeh et al. (2009). Amplification and sequencing of part of the ribosomal DNA (ITS region), translation elongation factor 1- $\alpha$  (EF1- $\alpha$ ) and  $\beta$ -tubulin genes were performed as described previously (Alves et al. 2006, Abdollahzadeh et al. 2009).

#### Phylogenetic analyses

Sequences were checked with BioEdit v. 7.0.9.0 (Hall 2006). The ITS and EF1- $\alpha$  sequences of two outgroups (*Neofusicoccum luteum* CBS 110299, CBS 110497) and additional isolates

were retrieved from GenBank. Sequences were aligned with ClustalX v. 1.83 (Thompson et al. 1997), using the following parameters: pairwise alignment parameters (gap opening = 10, gap extension = 0.1) and multiple alignment parameters (gap opening = 10, gap extension = 0.2, transition weight = 0.5, delay divergent sequences = 25 %). Alignments were checked and manual adjustments were made where necessary. Phylogenetic information contained in indels (gaps) was incorporated into the phylogenetic analyses using simple indel coding as implemented by GapCoder (Young & Healy 2003). Phylogenetic analyses were performed with PAUP v. 4.0b10 (Swofford 2003) for neighbour-joining (NJ) and maximum-parsimony (MP) analyses as described by Abdollahzadeh et al. (2010). A partition homogeneity test (PHT) was used to determine the congruence between the ITS, EF1- $\alpha$  and  $\beta$ -tubulin datasets (Farris et al. 1995, Huelsenbeck et al. 1996).

Bayesian analyses employing a Markov Chain Monte Carlo (MCMC) method were performed. The general time-reversible model of evolution (Rodriguez et al. 1990), including estimation of invariable sites and assuming a discrete gamma distribution with six rate categories (GTR+I+ $\Gamma$ ) was used. Four MCMC chains were run simultaneously, starting from random trees, for 10<sup>6</sup> generations. Trees were sampled every 100th generation for a total of 10<sup>4</sup> trees. The first 10<sup>3</sup> trees were discarded as the burn-in phase of each analysis. Posterior probabilities (Rannala & Yang 1996) were determined from a majority-rule consensus tree generated from the remaining 9 000 trees. The analysis was repeated three times starting from different random trees




Fig. 1 Distance tree obtained using the K2P substitution model on the combined ITS and EF1- $\alpha$  sequence data. BI/MP/NJ posterior probabilities and bootstrap support values are given at the nodes. The tree is rooted to *Neofusicoccum luteum* (CBS 110299, CBS 110497). New species and ex-type strains are in bold face.

to ensure trees from the same tree space were being sampled during each analysis. New sequences were deposited in GenBank (Table 1) and the alignment in TreeBASE (S14150).

## RESULTS

### Phylogenetic analyses

Phylogenetic analyses were done based on different combination of the three unlinked gene regions including ITS/EF1- $\alpha$ , ITS/ $\beta$ -tubulin and ITS/EF1- $\alpha$ / $\beta$ -tubulin. The phylogenies resulted from ITS/EF1- $\alpha$  were stable and reproducible with highly supported internal nodes, while  $\beta$ -tubulin did not improve the phylogenies so that combination of ITS and  $\beta$ -tubulin datasets resulted in poor and distinct phylogenies in different analyses (data not shown). Furthermore, PHT test for combined ITS/EF1- $\alpha$ / $\beta$ -tubulin datasets was significant ( $P < 0.01$ ). Therefore,  $\beta$ -tubulin was excluded and phylogenetic analyses were based on ITS and EF1- $\alpha$  sequences. The partition homogeneity test ( $P = 0.11$ ) indicated the phylogenies resulting from ITS and EF1- $\alpha$  were congruent so the ITS and EF1- $\alpha$  datasets were combined in a single analysis. The combined ITS and EF1- $\alpha$  sequences for 57 ingroup and 2 outgroup taxa contained 1 108 characters including alignment gaps, of which 334 characters were excluded, 528 were constant, 23 were variable and parsimony-uninformative and 223 were parsimony-informative. A heuristic search of the remaining 223 parsimony-informative characters resulted in a single most parsimonious tree of 426 steps (CI = 0.7, HI = 0.3, RI = 0.93). The Bayesian and NJ analyses produced phylogenetic trees with the same topology as the MP tree. The NJ tree is shown in Fig. 1 with BI/MP/NJ posterior probabilities and bootstrap support values at the nodes. MP and Bayesian trees are available in TreeBASE (S14150). In these phylogenetic analyses 22 subclades, representing 22 species of *Botryosphaeriaceae* with dark-walled 2-celled conidia, were recognized in two major clades corresponding to *Dothiorella* with 18 subclades, and *Spencermartinsia* with four subclades. Of these, nine subclades are recognised here as representatives of nine new species for the science.

### Taxonomy

All isolates in this study were induced to sporulate and produced pycnidia on poplar twigs on WA within 1–2 wk. No sexual structures were observed on the field specimens or in cultures. Based on ITS and EF1- $\alpha$  sequences, conidial and cultural characteristics and growth rate on MEA in the dark at 25 °C nine new species were identified. Of these, eight new species are described and illustrated here. Five species reside in the *Dothiorella* clade and the other three in the *Spencermartinsia* clade.

### Key to *Dothiorella* species<sup>1</sup>

1. Conidiomata papillate ..... 2
1. Conidiomata non-papillate ..... 8
2. Conidiomata with long necks (up to 1.5 mm) *D. longicollis*
2. Conidiomata with short necks (< 0.5 mm) ..... 3
3. Conidia striate ..... *D. striata*
3. Conidia smooth ..... 4
4. Conidial length 16–22  $\mu$ m ..... *D. dulcispiniae*
4. Conidial length > 21  $\mu$ m ..... 5
5. Colony growth rate on MEA in the dark at 25 °C > 30 mm/d (37 mm/d) ..... *D. prunicola*
5. Colony growth rate on MEA in the dark at 25 °C < 30 mm/d ..... 6

6. Colony growth rate on MEA in the dark at 25 °C > 20 mm/d ..... *D. pretoriensis*
6. Colony growth rate on MEA in the dark at 25 °C < 20 mm/d ..... 7
7. Colony growth rate on MEA in the dark at 25 °C < 15 mm (14 mm/d) ..... *D. iranica*
7. Colony growth rate on MEA in the dark at 25 °C > 15 mm (17.9 mm/d) ..... *D. brevicollis*
8. Conidial length < 16  $\mu$ m (av. length 15  $\mu$ m) *D. americana*
8. Conidial length 16  $\mu$ m or more (av. length > 18  $\mu$ m) ... 9
9. Average conidial width > 10  $\mu$ m ..... 10
9. Average conidial width < 10  $\mu$ m ..... 13
10. Average conidial length < 21  $\mu$ m ..... 11
10. Average conidial length > 21  $\mu$ m ..... 12
11. Average conidial width > 11  $\mu$ m, conidial L/W ratio 1.7, colonies reaching 25–30 mm on MEA after 4 d in the dark at 25 °C ..... *D. parva*
11. Average conidial width < 11  $\mu$ m, conidial L/W ratio 2, colonies reaching 50–70 mm on MEA after 4 d in the dark at 25 °C ..... *D. sempervirentis*
12. Conidia 23–31  $\times$  9–11  $\mu$ m (av. 27.1  $\times$  10.8  $\mu$ m) ..... *D. casuarini*
12. Conidia 23–23.4  $\times$  10.8–11  $\mu$ m (av. 23.2  $\times$  10.9  $\mu$ m) ... *D. iberica*
13. Average conidial length < 20  $\mu$ m ..... 14
13. Average conidial length > 20  $\mu$ m ..... 15
14. Conidial L/W ratio 2 ..... *D. santali*
14. Conidial L/W ratio 2.4 ..... *D. moneti*
15. Conidia 21.4–21.9  $\times$  9.7–9.9  $\mu$ m (L/W ratio 2.2) ..... *D. sarmorum*<sup>2</sup>
15. Conidia 22–22.5  $\times$  9–9.5  $\mu$ m (L/W ratio 2.4) ..... *D. uruguayensis*<sup>2</sup>

***Dothiorella iranica*** Abdollahz. & A.J.L. Phillips, *sp. nov.* — MycoBank MB803988; Fig. 2

*Etymology.* Named for the country of origin, Iran.

*Conidiomata* pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 370  $\mu$ m diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, papillate with a central ostiole. *Conidiophores* absent. *Conidiogenous cells* cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth, (6.5–)9–12(–18.4)  $\times$  2–5  $\mu$ m. *Conidia* subcylindrical to ellipsoid or ovoid, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends rounded, often with a truncate base, rarely becoming tapered at the base, (21.9–)23–26(–27.5)  $\times$  (8.8–)9–11(–11.2)  $\mu$ m, 95 % confidence limits = 25–25.7  $\times$  9.9–10.2  $\mu$ m (av.  $\pm$  S.D. = 25.3  $\pm$  1.4  $\times$  10.1  $\pm$  0.6  $\mu$ m, l/w ratio = 2.5  $\pm$  0.2).

*Culture characteristics* — Colonies cottony with aerial mycelium, aerial mycelium becoming olivaceous-buff to grey-olivaceous at the surface and isabelline to dull green at the reverse after 2 wk in the dark at 25 °C. Colonies reaching 56 mm on MEA after 4 d in the dark at 25 °C. Cardinal temperatures for growth: min.  $\leq$  5 °C, max.  $\geq$  35 °C, opt. 25 °C.


*Substrate* — Twigs of *Olea europaea*.

*Distribution* — Northern Iran.

*Specimens examined.* IRAN. Golestan Province, Gorgan (Agriculture Research Center), on twigs of *Olea europaea*, June 2007, A. Javadi, holotype IRAN 16253F, culture ex-type IRAN 1587C = CBS 124722.

<sup>1</sup> This key is based on conidial morphology and culture characteristics.

<sup>2</sup> These two species are phylogenetically distant.



**Fig. 2** *Dothiorella iranica* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline immature conidia developing on conidiogenous cells; d, e. brown aseptate conidia on conidiogenous cells; f. brown 1-septate conidia attached to the conidiogenous cells; g. mature conidia; h. mature conidia in two different focal planes. — Scale bars: a = 250  $\mu$ m; b, d–g = 10  $\mu$ m; c, h = 5  $\mu$ m.

**Notes** — Phylogenetically this species resides in a distinct subclade in *Dothiorella* and morphologically conidia of *D. iranica* are longer ( $25.3 \pm 1.4 \times 10.1 \pm 0.6 \mu\text{m}$ , l/w ratio =  $2.5 \pm 0.2$ ) than those of all other *Dothiorella* spp., except *D. casuarini* ( $27.1 \times 10.8$ ).

***Dothiorella parva*** Abdollahz., Zare & A.J.L. Phillips, sp. nov. — MycoBank MB803989; Fig. 3

**Etymology.** Named for its short conidia.

**Conidiomata** pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 350  $\mu$ m diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, non-papillate with a central ostiole. **Conidiophores** absent. **Conidiogenous cells** cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth, (7–)8–11(–13.8)


$\times 3\text{--}5 \mu\text{m}$ . **Conidia** ellipsoid to ovoid, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends rounded, often with a truncate base, (17.2–)18–21(–23.8)  $\times$  (8.9–)10–13(–15.1)  $\mu\text{m}$ , 95 % confidence limits =  $19.4\text{--}19.8 \times 11.4\text{--}12$  (av.  $\pm$  S.D. =  $19.6 \pm 1.2 \times 11.7 \pm 1.6 \mu\text{m}$ , l/w ratio =  $1.7 \pm 0.2$ ).

**Culture characteristics** — Colonies cottony with aerial mycelium, aerial mycelium becoming pale olivaceous-grey to iron-grey at the surface and olivaceous-buff to dull green at the reverse after 2 wk in the dark at 25  $^{\circ}\text{C}$ . Colonies reaching 25–30 mm on MEA after 4 d in the dark at 25  $^{\circ}\text{C}$ . Cardinal temperatures for growth: min.  $\leq 5 \text{ }^{\circ}\text{C}$ , max.  $\geq 35 \text{ }^{\circ}\text{C}$ , opt. 25  $^{\circ}\text{C}$ .

**Substrate** — Twigs of *Corylus avellana*.

**Distribution** — North-west of Iran.

**Specimens examined.** IRAN, Ardabil Province, Ardabil (Fandoghlo Forest Park), on twigs of *Corylus avellana*, July 2007, J. Abdollahzadeh and A. Javadi, holotype IRAN 14264F, culture ex-type IRAN 1579C = CBS 124720. Additional isolates are given in Table 1.



**Fig. 3** *Dothiorella parva* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline immature conidia developing on conidiogenous cells; d. brown aseptate and 1-septate conidia attached to the conidiogenous cells; e. mature conidia; f. microconidiogenous cells; g. microconidia. — Scale bars: a = 1 000  $\mu$ m; b, c, f, g = 5  $\mu$ m; d, e = 10  $\mu$ m.

Notes — Phylogenetically, *D. parva* is closely related to *D. sempervirens* and *Dothiorella* sp., but can be distinguished from these two species on account of its shorter and wider conidia ( $19.6 \pm 1.2 \times 11.7 \pm 1.6 \mu\text{m}$ , l/w ratio =  $1.7 \pm 0.2$ ) and slower growth rate on MEA in the dark at 25 °C. This species differed in nucleotide sequences from *D. sempervirens* (three substitutions in ITS, five substitutions and 2 insertions/deletions in EF1- $\alpha$ ) and *Dothiorella* sp. (one substitution in ITS, five substitutions and one insertion/deletion in EF1- $\alpha$ ).

***Dothiorella prunicola*** A.J.L. Phillips & Abdollahz., *sp. nov.* — MycoBank MB803991; Fig. 4

*Etymology.* Named for the host it was first isolated from, namely *Prunus*.

*Conidiomata* pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 370  $\mu\text{m}$  diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, papillate with a central ostiole. *Conidiophores* absent. *Conidiogenous cells* cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth,  $(5.9\text{--})9\text{--}14\text{--}(20) \times 3\text{--}5 \mu\text{m}$ . *Conidia* subcylindrical to ellipsoid or ovoid, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends rounded, often with a truncate base,  $(19\text{--})22\text{--}27\text{--}(30.5) \times (10.5\text{--})11\text{--}14\text{--}(16.8) \mu\text{m}$ , 95 % confidence limits =  $23.9\text{--}25.1 \times 12.5\text{--}13.1 \mu\text{m}$  (av.  $\pm$  S.D. =  $24.5 \pm 2.3 \times 12.8 \pm 1.2 \mu\text{m}$ , l/w ratio =  $1.9 \pm 0.2$ ). *Spermatogenous cells* discrete or integrated, hyaline, smooth, cylindrical, holoblastic or proliferating via phialides with periclinal thickenings,  $5.8\text{--}17.3 \times 1\text{--}3 \mu\text{m}$ . *Spermatia* hyaline, smooth, aseptate, rod-shaped with rounded ends,  $3.5\text{--}5.3 \times 1\text{--}2 \mu\text{m}$ .

*Culture characteristics* — Colonies rosette with lobed margins and aerial mycelium, white to cream at the surface and reverse after 2 wk in the dark at 25 °C. Colonies reaching 37 mm on MEA after 4 d in the dark at 25 °C. Cardinal temperatures for growth: min.  $\leq 5 \text{ }^\circ\text{C}$ , max.  $\geq 35 \text{ }^\circ\text{C}$ , opt. 25 °C.

*Substrate* — Twigs of *Prunus dulcis*.

*Distribution* — Portugal.

*Specimens examined.* PORTUGAL, Algarve Province, on twigs of *Prunus dulcis*, June 2007, E. Diogo, holotype IRAN 16252F, culture ex-type IRAN 1541C = CBS 124723.

Notes — This species resides in a completely distinct sub-clade within *Dothiorella*. Morphologically, *D. prunicola* resem-

bles *D. brevicollis*, *D. iranica* and *D. pretoriensis* but can be distinguished on average conidial dimensions ( $24.5 \pm 2.3 \times 12.8 \pm 1.2 \mu\text{m}$ , l/w ratio =  $1.9 \pm 0.2$ ) and growth rate on MEA in the dark at 25 °C. Moreover, *D. prunicola* can be distinguished from all *Dothiorella* spp. on account of its white and creamy colony colour, a feature that is never seen in the family *Botryosphaeriaceae*.

***Dothiorella sempervirens*** Abdollahz., Zare & A.J.L. Phillips, *sp. nov.* — MycoBank MB803987; Fig. 5

*Etymology.* Named for the host species it was first isolated from.

*Conidiomata* pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 510  $\mu\text{m}$  diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, non-papillate with a central ostiole. *Conidiophores* absent. *Conidiogenous cells* cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings or rarely proliferating percurrently to form one or two annellations, hyaline, thin-walled, smooth,  $(8.2\text{--})9\text{--}11\text{--}(17.3) \times 3\text{--}5 \mu\text{m}$ . *Conidia* subcylindrical to ellipsoid or ovoid, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends rounded, often with a truncate base,  $(16.1\text{--})18\text{--}20\text{--}(22.8) \times (8.2\text{--})9\text{--}11\text{--}(11.7) \mu\text{m}$ , 95 % confidence limits =  $19.8\text{--}20.3 \times 10\text{--}10.4 \mu\text{m}$  (av.  $\pm$  S.D. =  $20.1 \pm 1.3 \times 10.2 \pm 0.9 \mu\text{m}$ , l/w ratio =  $2 \pm 0.2$ ).


*Culture characteristics* — Colonies appressed with a sparse aerial mycelium at the margin, grey-olivaceous to olivaceous-black at the surface and greenish olivaceous to dull green in reverse after 2 wk in the dark at 25 °C. Colonies reaching 50–70 mm on MEA after 4 d in the dark at 25 °C. Cardinal temperatures for growth: min.  $\leq 5 \text{ }^\circ\text{C}$ , max.  $\geq 35 \text{ }^\circ\text{C}$ , opt. 25 °C.

*Substrate* — Twigs and cones of *Cupressus sempervirens*.


*Distribution* — Northern Iran.

*Specimens examined.* IRAN, Golestan Province, Gorgan (City Park), on twigs and cones of *Cupressus sempervirens*, Aug. 2006, M.A. Aghajani, holotype IRAN 14265F, culture ex-type IRAN 1583C = CBS 124718. Additional isolates are given in Table 1.

Notes — Phylogenetically, *D. sempervirens* is closely related to *D. parva* and *Dothiorella* sp., but can be distinguished on average conidial dimensions ( $20.1 \pm 1.3 \times 10.2 \pm 0.9 \mu\text{m}$ , l/w ratio =  $2 \pm 0.2$ ) and growth rate on MEA in the dark at 25 °C. Compared with *Dothiorella* sp. nine differences were detected



**Fig. 4** *Dothiorella prunicola* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline immature conidia developing on conidiogenous cells; d. brown 1-septate conidia attached to the conidiogenous cells; e. mature conidia; f. microconidiogenous cells; g. microconidia. — Scale bars: a = 250  $\mu\text{m}$ ; b, c, f, g = 5  $\mu\text{m}$ ; d, e = 10  $\mu\text{m}$ .



**Fig. 5** *Dothiorella sempervirens* holotype. a. Conidiomata on poplar twigs in culture; b–d. hyaline immature conidia developing on conidiogenous cells; e, f. brown aseptate and 1-septate conidia attached to the conidiogenous cells; g. mature conidia. — Scale bars: a = 1 000 µm; b, e–g = 10 µm; c, d = 5 µm.

in ITS and EF1-α sequences (four substitutions in ITS, four substitutions and one insertion/deletion in EF1-α).

***Dothiorella striata*** A.J.L. Phillips & Abdollahz., *sp. nov.* — MycoBank MB803990; Fig. 6

*Etymology.* Named for the distinctive striations on the conidial wall.

**Conidiomata** pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 420 µm diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, papillate with a central ostiole. **Conidiophores** absent. **Conidiogenous cells** cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth, (5.9–)9–14(–20) × 3–5 µm. **Conidia** subcylindrical to ellipsoid or ovoid, brown, 1-septate, striate, thumb-like striations visible on hyaline conidia even while attached to conidiogenous cells, occasionally slightly constricted at septum, moderately thick-walled, internally finely verruculose, ends rounded, often with a truncate base, (21–)23–26(–29.4) × (8.9–)9–12(–15.1) µm, 95 % confidence limits = 24.9–25.4 × 10.5–11 µm (av. ± S.D. = 25.1 ± 1.4 × 10.7 ± 1.2 µm, l/w ratio = 2.4 ± 0.3).

**Culture characteristics** — Colonies with abundant aerial mycelia reaching the lid of Petri plates, aerial mycelium becoming smoke-grey to olivaceous-black at the surface and greenish olivaceous to dull green at the reverse after 2 wk in the dark at 25 °C. Colonies reaching 45–55 mm on MEA after 4 d in the dark at 25 °C. Cardinal temperatures for growth: min. ≤ 5 °C, max. ≥ 35 °C, opt. 25 °C.

**Substrate** — Twigs of *Citrus sinensis*.

**Distribution** — New Zealand.

**Specimens examined.** NEW ZEALAND, Northland, Kerikeri, Collins Orchard, Inlet Road, on twigs of *C. sinensis*, Sept. 2006, S.R. Pennycook, P.R. Johnston and B.C. Paulus, holotype PDD 92029, culture ex-type ICMP 16824 = CBS 124731. Additional isolates are given in Table 1.

**Notes** — Phylogenetically, *D. striata* is closely related to *D. uruguayensis*, but it can be distinguished from all *Dothiorella* spp. on account of the striate conidia, which was never

seen in any other *Dothiorella* species. Furthermore, *D. striata* differs from all other species on account of colonies with abundant aerial mycelia reaching the lid of Petri plates.

***Dothiorella* sp.**

**Notes** — Phylogenetically, *Dothiorella* sp. is closely related to *D. parva* and *D. sempervirens*, but can be distinguished on account of the number of differences in ITS and EF1-α sequences, longer and narrower conidia and faster growth rate on MEA in the dark at 25 °C. This species is being further studied in relation to other species collected from *Corylus* and therefore is not formally described here.

**Key to *Spencermartinsia* species<sup>1</sup>**


- 1. Average conidial length < 20 µm . . . . . *S. mangiferae*
- 1. Average conidial length > 20 µm . . . . . 2
- 2. Conidia truncate at both ends (av. 25.8 × 12.2 µm) . . . . .
- . . . . . *S. citricola*
- 2. Conidia round at apex, often with a truncate base . . . . . 3
- 3. Average conidial width > 10 µm (av. 22.5 × 11 µm) . . . . .
- . . . . . *S. plurivora*
- 3. Average conidial width < 10 µm (av. 20.4 × 9.3 µm) . . . . .
- . . . . . *S. viticola*

***Spencermartinsia citricola*** A.J.L. Phillips & Abdollahz., *sp. nov.* — MycoBank MB803992; Fig. 7

*Etymology.* Named for the host it was first isolated from, namely *Citrus*.

**Conidiomata** pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 460 µm diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, non-papillate with a central ostiole. **Conidiophores** absent. **Conidiogenous cells** cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth, (7.6–)9–11(–12)


<sup>1</sup> This key is based on conidial morphology.



**Fig. 6** *Dothiorella striata* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline immature conidia developing on conidiogenous cells; d. brown aseptate and 1-septate conidia attached to the conidiogenous cells; e. hyaline and brown striate conidia attached to the conidiogenous cells; f. mature conidia; g. mature conidia with striation. — Scale bars: a = 1 000  $\mu\text{m}$ ; b, d, e = 5  $\mu\text{m}$ ; c, f, g = 10  $\mu\text{m}$ .

$\times 2\text{--}4$   $\mu\text{m}$ . *Conidia* oblong to subcylindrical, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends truncate,  $(23.7\text{--})24\text{--}27(\text{--}28) \times (9.5\text{--})10\text{--}12(\text{--}14.1)$   $\mu\text{m}$ , 95 % confidence limits =  $25.5\text{--}26 \times 11.9\text{--}12.5$   $\mu\text{m}$  (av.  $\pm$  S.D. =  $25.8 \pm 1.1 \times 12.2 \pm 1.3$   $\mu\text{m}$ , l/w ratio =  $2.1 \pm 0.2$ ).

**Culture characteristics** — Colonies cottony with dense aerial mycelium and crenate margins, aerial mycelium becoming smoke-grey to olivaceous-black at the surface and greenish olivaceous to dull green at the reverse after 2 wk in the dark at 25  $^{\circ}\text{C}$ . Colonies reaching 15–20 mm on MEA after 4 d in the dark at 25  $^{\circ}\text{C}$ . Cardinal temperatures for growth: min.  $\leq 5$   $^{\circ}\text{C}$ , max.  $\geq 35$   $^{\circ}\text{C}$ , opt. 25  $^{\circ}\text{C}$ .



**Fig. 7** *Spencermartinsia citricola* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline immature conidia developing on conidiogenous cells; d, e. brown aseptate and 1-septate conidia attached to the conidiogenous cells; f. mature conidia. — Scale bars: a = 1 000  $\mu\text{m}$ ; b–e = 5  $\mu\text{m}$ ; f = 10  $\mu\text{m}$ .

Substrate — Twigs of *Citrus sinensis*.  
 Distribution — New Zealand.

*Specimens examined.* NEW ZEALAND, Northland, Kerikeri, Collins Orchard, Inlet Road, on twigs of *C. sinensis*, Sept. 2006, S.R. Pennycook, P.R. Johnston and B.C. Paulus, holotype PDD92023, culture ex-type ICMP16828 = CBS124729. Additional isolates are given in Table 1.

*Notes* — Phylogenetically, *S. citricola* is closely related to *S. mangiferae* and *S. plurivora*, but morphologically can be distinguished an account of conidial dimensions ( $25.8 \pm 1.1 \times 12.2 \pm 1.3 \mu\text{m}$ ), shape (truncated at either end) and slower growth rate on MEA in the dark at 25 °C. This species differed in nucleotide sequences from *S. mangiferae* (six substitutions and 2 insertions/deletions in EF1- $\alpha$ ) and *S. plurivora* (two substitutions in ITS, six substitutions and three insertions/deletions in EF1- $\alpha$ ).

***Spencermartinsia mangiferae*** Abdollahz., Javadi & A.J.L. Phillips, *sp. nov.* — MycoBank MB803993; Fig. 8

*Etymology.* Named for the host it was first isolated from, namely *Mangifera*.

*Conidiomata* pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 400  $\mu\text{m}$  diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, non-papillate with a central ostiole. *Conidiophores* absent. *Conidiogenous cells* cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth, (5.2–)6–9(–11.8)  $\times$  3–5  $\mu\text{m}$ . *Conidia* subcylindrical to ellipsoid or ovoid, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends rounded, often with a truncate base, (14.4–)17–22(–22.5)  $\times$  (6.3–)8–10(–11)  $\mu\text{m}$ , 95 % confidence limits = 18.8–19.2  $\times$  8.9–9.1  $\mu\text{m}$  (av.  $\pm$  S.D. =  $19 \pm 1.6 \times 9 \pm 0.9 \mu\text{m}$ , l/w ratio =  $2.1 \pm 0.2$ ).

*Culture characteristics* — Colonies cottony with dense aerial mycelium, aerial mycelium becoming smoke-grey to olivaceous-grey at the surface and grey-olivaceous to olivaceous-black at the reverse after 2 wk in the dark at 25 °C. Colonies reaching

50–85 mm on MEA after 4 d in the dark at 25 °C. Cardinal temperatures for growth: min.  $\leq 5 \text{ }^\circ\text{C}$ , max.  $\geq 35 \text{ }^\circ\text{C}$ , opt. 25 °C.

Substrate — Twigs of *Mangifera indica*.  
 Distribution — Southern Iran.

*Specimens examined.* IRAN, Hormozgan Province, Bandar Abbas (Hajjabad-Siaho), on twigs of *M. indica*, Mar. 2007, J. Abdollahzadeh and A. Javadi, holotype IRAN 14266F, culture ex-type IRAN 1584C = CBS 124727.


*Notes* — Phylogenetically, *S. mangiferae* is closely related to *S. citricola* and *S. plurivora*, but in terms of morphology it is differed from all other species by having small conidia ( $19 \pm 1.6 \times 9 \pm 0.9 \mu\text{m}$ ). This species also differed in nucleotide sequences from *S. citricola* (six substitutions and two insertions/deletions in EF1- $\alpha$ ) and *S. plurivora* (two substitutions in ITS, six substitutions and one insertion/deletion in EF1- $\alpha$ ).

***Spencermartinsia plurivora*** Abdollahz., Javadi & A.J.L. Phillips, *sp. nov.* — MycoBank MB803994; Fig. 9


*Etymology.* Named for its broad host range.

*Conidiomata* pycnidial, produced on poplar twigs on WA within 1–2 wk, solitary or aggregated, individual conidiomata globose, up to 420  $\mu\text{m}$  diam, superficial or semi-immersed, covered with hyphal hairs, uniloculate, thick-walled, non-papillate with a central ostiole. *Conidiophores* absent. *Conidiogenous cells* cylindrical to lageniform, discrete or integrated, holoblastic, indeterminate, proliferating at the same level giving rise to periclinal thickenings, hyaline, thin-walled, smooth, (5.1–)7–10(–11.9)  $\times$  3–5  $\mu\text{m}$ . *Conidia* subcylindrical to ellipsoid or ovoid, brown, 1-septate, occasionally slightly constricted at septum, moderately thick-walled, externally smooth, internally finely verruculose, ends rounded, often with a truncate base, (18–)20–25(–27)  $\times$  (8.9–)10–13(–14.4)  $\mu\text{m}$ , 95 % confidence limits = 22.3–22.7  $\times$  10.8–11.2  $\mu\text{m}$  (av.  $\pm$  S.D. =  $22.5 \pm 1.7 \times 11 \pm 1.1 \mu\text{m}$ , l/w ratio =  $2.1 \pm 0.2$ ).

*Culture characteristics* — Colonies cottony with dense aerial mycelium, aerial mycelium becoming smoke-grey to olivaceous-grey at the surface and grey-olivaceous to olivaceous-black at the reverse after 2 wk in the dark at 25 °C. Colonies reaching 62–84 mm on MEA after 4 d in the dark at 25 °C. Cardinal temperatures for growth: min.  $\leq 5 \text{ }^\circ\text{C}$ , max.  $\geq 35 \text{ }^\circ\text{C}$ , opt. 25 °C.



**Fig. 8** *Spencermartinsia mangiferae* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline immature conidia developing on conidiogenous cells; d. brown aseptate conidia on conidiogenous cells; e, f. mature conidia in two different focal planes. — Scale bars: a = 1 000  $\mu\text{m}$ ; b, d, e = 10  $\mu\text{m}$ ; c, f = 5  $\mu\text{m}$ .



**Fig. 9** *Spencermartinsia plurivora* holotype. a. Conidiomata on poplar twigs in culture; b, c. hyaline and immature conidia developing on conidiogenous cells; d, e. brown aseptate and 1-septate conidia attached to the conidiogenous cells; f. mature conidia. — Scale bars: a = 1 000  $\mu$ m; b, c = 5  $\mu$ m; d–f = 10  $\mu$ m.

**Substrate** — Twigs of *Casuarina* sp., *Citrus* sp., *Cupressus sempervirens*, *Eucalyptus* sp., *Juglans regia*, *Malus domestica*, *Prunus armeniaca* and *Vitis vinifera*.

**Distribution** — Southern Iran, Spain.

**Specimens examined.** IRAN, Khuzestan Province, Dezful (Safiabad Citrus Research Centre), on twigs of *Citrus* sp., Nov. 2006, J. Abdollahzadeh and A. Javadi, holotype IRAN 14267F, culture ex-type IRAN 1557C = CBS 124724.

**Notes** — Phylogenetically, *S. plurivora* is closely related to *S. citricola* and *S. mangiferae*, but morphologically it resembles *S. viticola*. This species can be distinguished from other species on account of its conidial dimensions ( $22.5 \pm 1.7 \times 11 \pm 1.1 \mu$ m) and colony growth rate on MEA at 25 °C. Moreover, this species differed in nucleotide sequences from *S. citricola* (two substitutions in ITS, six substitutions and three insertions/deletions in EF1- $\alpha$ ) and *S. mangiferae* (two substitutions in ITS, six substitutions and one insertion/deletion in EF1- $\alpha$ ).

## DISCUSSION

In this study the phylogenetic analyses based on ITS and EF1- $\alpha$  sequences revealed two major clades corresponding to *Dothiorella* and *Spencermartinsia*. Within *Dothiorella* 18 subclades were resolved. Species names are available for 11 of these subclades, and five new species are introduced here from isolates collected from different woody hosts in Iran, New Zealand, Portugal and Spain. We refrain from giving species names to the subclade bearing *Diplodia juglandis*, *Diplodia coryli* and the two isolates IRAN 1570C and IRAN 1573C, since this will be dealt with in a separate paper currently being prepared on the species of *Dothiorella* on *Corylus*. The subclade bearing two isolates CMW 25404 and CMW 25405 was recognised by Jami et al. (2012) as *Spencermartinsia* sp., but they have not described it thus far. Four subclades were resolved in *Spencermartinsia* representing four species; *S. viticola* and three species which are newly described in this paper.

*Spencermartinsia* was introduced by Phillips et al. (2008) on account of its phylogenetic distinction from *Dothiorella*. The presence of apiculi on the ascospores of *Spencermartinsia viticola* was the only morphological character that separates it from *Dothiorella*, the ascospores of which do not have apiculi.

However, a sexual state is known only for *S. viticola*, and there are no distinctive asexual characters that differentiate these two genera.

All *Dothiorella* species can be distinguished based on ITS and EF1- $\alpha$  sequence data supported with conidial dimensions (Table 2). In this survey, we also considered growth rate, a physiological character, on MEA in the dark at 25 °C to differentiate some closely related species, and apparently this is a useful character in some cases. *Dothiorella striata* was isolated from twigs of *Citrus* in New Zealand. In a phylogenetic study based on SSU, LSU, ITS,  $\beta$ -tubulin and EF1- $\alpha$  sequences the two isolates representing *D. striata* formed a distinct clade as a sister group to *Spencermartinsia* with quite low support in MP analysis (Phillips et al. 2008). The presence of striate conidia as a strong and unique morphological character led Phillips et al. (2008) to suspect that this clade could represent a separate genus. However, they declined to introduce a new genus and did not describe this species because the isolates failed to sporulate. In the present study isolates of *D. striata* clearly lie within the *Dothiorella* clade and cultures sporulating on poplar twigs confirmed the presence of conidial striation in this species. Therefore, conidial striation is interpreted as a distinctive morphological character at the species level in the *Dothiorella* clade.

Thus far, 18 species of *Dothiorella*, including those identified in this study, have been characterised from different hosts based on combined morphology and DNA sequence data (Luque et al. 2005, Phillips et al. 2005, Pavlic et al. 2008, Taylor et al. 2009, de Wet et al. 2009, Urbez-Torres et al. 2011, Jami et al. 2012). Of these, *D. sarmentorum* is cosmopolitan and has been isolated from 34 different host species across six continents, and *D. iberica* has been found on seven different tree species in Algeria, Italy, Spain and USA (<http://nt.ars-grin.gov/fungal-databases>). However, other species have a very narrow host range and limited geographic distribution. To obtain a more realistic conclusion on host ranges, we require more sampling from various hosts in relatively unexplored regions.

*Spencermartinsia* was introduced as a monotypic genus typified with *S. viticola*, a species reported from four different woody hosts (mainly *Vitis vinifera*) in China, South Africa, Spain, USA

**Table 2** Conidial dimensions of *Dothiorella* and *Spencermartinsia* species investigated in this study and previous studies.

Species	Conidial dimensions ( $\mu\text{m}$ )	Average ( $\mu\text{m}$ )	L/W ratio	Reference
<i>D. americana</i>	14.2–15.8 $\times$ 5.7–6.6	15 $\times$ 6.1	2.4	Urbez Torres et al. 2012
<i>D. brevicollis</i>	21.5–26 $\times$ 9–12	–	–	Jami et al. 2012
<i>D. casuarini</i>	23–31 $\times$ 9–12	27.1 $\times$ 10.8	–	de Wet et al. 2009
<i>D. dulcispinae</i>	16–22 $\times$ 7–10	–	–	Jami et al. 2012
<i>D. iberica</i>	23–23.4 $\times$ 10.8–11	23.2 $\times$ 10.9	2.2	Phillips et al. 2005
<i>D. iranica</i>	23–26 $\times$ 9–11	25.3 $\times$ 10.1	2.5	This study
<i>D. longicollis</i>	19–22 $\times$ 8–9.5	20.4 $\times$ 8.7	2.3	Pavlic et al. 2008
<i>D. moneti</i>	17–22 $\times$ 7–10	19.8 $\times$ 8.4	2.4	Taylor et al. 2009
<i>D. parva</i>	18–21 $\times$ 10–13	19.6 $\times$ 11.7	1.7	This study
<i>D. pretoriensis</i>	20–28 $\times$ 7–14	–	–	Jami et al. 2012
<i>D. prunicola</i>	22–27 $\times$ 11–14	24.5 $\times$ 12.8	1.9	This study
<i>D. santali</i>	16–20 $\times$ 7–11	18.2 $\times$ 9	2	Taylor et al. 2009
<i>D. sarmentorum</i>	21.4–21.9 $\times$ 9.7–9.9	21.6 $\times$ 9.8	2.2	Phillips et al. 2005
<i>D. sempervirentis</i>	18–20 $\times$ 9–11	20.1 $\times$ 10.2	2	This study
<i>D. striata</i>	23–26 $\times$ 9–12	25.1 $\times$ 10.7	2.4	This study
<i>D. uruguayensis</i>	22–22.5 $\times$ 9–9.5	–	–	Pérez et al. 2010
<i>Dothiorella</i> sp.	21–27 $\times$ 8–10	24 $\times$ 9.9	2.2	This study
<i>S. citricola</i>	24–27 $\times$ 10–12	25.8 $\times$ 12.2	2.1	This study
<i>S. mangiferae</i>	17–22 $\times$ 8–10	19 $\times$ 9	2.1	This study
<i>S. plurivora</i>	20–25 $\times$ 10–13	22.5 $\times$ 11	2.1	This study
<i>S. viticola</i>	20.2–20.6 $\times$ 9.2–9.4	20.4 $\times$ 9.3	2.2	Luque et al. 2005

and Uruguay (<http://nt.ars-grin.gov/fungaldatabases>). Based on an ITS and EF1- $\alpha$  phylogeny, Phillips et al. (2013) transferred the two recently described species, *S. uruguayensis* and *S. pretoriensis* to *Dothiorella*, and introduced two new combinations: *D. uruguayensis* and *D. pretoriensis*. In the present phylogenetic study we introduce a further three species. Therefore, only four species remain in *Spencermartinsia*. These species can be differentiated based on conidial dimensions (Table 2) and shape. As discussed in the case of *Dothiorella*, growth rate on MEA in the dark at 25 °C is obviously a helpful character to separate some closely related species in *Spencermartinsia*, as *S. citricola* is distinct from two closely related species, *S. plurivora* and *S. mangiferae*, with a much slower growth rate. According to Phillips et al. (2008) the isolate (CBS 117006) collected from *V. vinifera* in Spain was phylogenetically separate from *S. viticola* and produced a red-brown pigment. In the present study this isolate grouped with Iranian isolates from different woody hosts in a distinct clade corresponding to a new species we have named *S. plurivora*. However, in this study none of the Iranian isolates produced any pigment, which is consistent with the observations of Abdollahzadeh et al. (2010, 2013) about the limited taxonomic value of cultural characteristics in differentiation of *Botryosphaeriaceae* species. Furthermore, *S. plurivora*, the most common species, was characterised on eight different woody hosts in Iran (14 isolates) and Spain (1 isolate), while *S. mangiferae* was found on mango in Iran and *S. citricola* on citrus in New Zealand.

Although pathogenicity of the species described in this study has not been determined, according to pathogenicity tests previously conducted by different researchers (van Niekerk et al. 2004, Luque et al. 2005, Damm et al. 2007, Taylor et al. 2009, Inderbitzin et al. 2010, Urbez-Torres et al. 2012), the species of *Dothiorella* and *Spencermartinsia* appear to be minor pathogens or can be considered as saprophytic or endophytic fungi in association with different woody plants. Pathogenicity, host specificity and geographic distribution of the characterised species remain unknown issues that should be considered in future studies.

**Acknowledgements** Part of this work was financed by the European Regional Development Fund and Fundação para a Ciência e a Tecnologia (FCT) Portugal under project PPCDT/AGR/56140/2004, and through grant PEST-OE/BIA/UI0457/2011. A.J.L. Phillips was supported by grant number SFRH/BCC/15810/2005 from FCT, and J. Abdollahzadeh received a grant from Studienstiftung Mykologie, Köln, Germany and Kurdistan provincial office.

Shaun Pennycook and Peter Johnston, Landcare Research, Auckland, New Zealand provided isolates, type specimens and culture collection numbers for *D. striata* and *S. citricola*. M.A. Aghajani, Agricultural and Natural Resources Research Center of Golestan Province, Iran provided *D. sempervirentis* samples.

## REFERENCES

- Abdollahzadeh J, Javadi A, Mohammadi Goltapeh E, Zare R, Phillips AJL. 2010. Phylogeny and morphology of four new species of Lasiodiplodia from Iran. *Persoonia* 25: 1–10.
- Abdollahzadeh J, Mohammadi Goltapeh E, Javadi A, Shams-bakhsh M, Zare R, Phillips AJL. 2009. Barriopsis iraniana and Phaeobotryon cupressi: two new species of the Botryosphaeriaceae from trees in Iran. *Persoonia* 23: 1–8.
- Abdollahzadeh J, Zare R, Phillips AJL. 2013. Phylogeny and taxonomy of Botryosphaeria and Neofusicoccum species in Iran, with description of Botryosphaeria scharifii sp. nov. *Mycologia* 105: 220–230.
- Alves A, Correia A, Phillips AJL. 2006. Multi-gene genealogies and morphological data support Diplodia cupressi sp. nov., previously recognized as D. pinea f. sp. cupressi, as a distinct species. *Fungal Diversity* 23: 1–15.
- Crous PW, Palm ME. 1999. Reassessment of the anamorph genera Botryodiplodia, Dothiorella and Fusicoccum. *Sydowia* 51: 161–175.
- Damm U, Crous PW, Fourie PH. 2007. Botryosphaeriaceae as potential pathogens of Prunus species in South Africa, with descriptions of Diplodia africana and Lasiodiplodia plurivora sp. nov. *Mycologia* 99: 664–680.
- Denman S, Crous PW, Taylor JE, Kang JC, Pascoe I, Wingfield MJ. 2000. An overview of the taxonomic history of Botryosphaeria and a re-evaluation of its anamorphs based on morphology and ITS rDNA phylogeny. *Studies in Mycology* 45: 129–140.
- Farris JS, Kallersjö M, Kluge AG, Bult C. 1995. Testing significance of incongruence. *Cladistics* 10: 315–319.
- Hall T. 2006. Bioedit 7.5.0.3. Department of Microbiology, North Carolina State University. <http://www.mbio.ncsu.edu/BioEdit/Bioedit.html>.
- Huelsenbeck JP, Bull JJ, Cunningham CV. 1996. Combining data in phylogenetic analysis. *Trends in Ecology & Evolution* 11: 152–158.
- Inderbitzin P, Bostock RM, Trouillas FP, Michailides TJ. 2010. A six locus phylogeny reveals high species diversity in Botryosphaeriaceae from California almond. *Mycologia* 102: 1350–1368.
- Jami F, Slippers B, Wingfield MJ, Gryzenhout M. 2012. Five new species of the Botryosphaeriaceae from Acacia karroo in South Africa. *Cryptogamie Mycologie* 33: 245–266.
- Liu JK, Phookamsak R, Doilom M, Wikee S, Li YM, et al. 2012. Towards a natural classification of Botryosphaeriales. *Fungal Diversity* 57: 149–210.
- Luque J, Martos S, Phillips AJL. 2005. Botryosphaeria viticola sp. nov. on grapevines: a new species with a Dothiorella anamorph. *Mycologia* 97: 1111–1121.
- Niekerk JM van, Crous PW, Groenewald JZ, Fourie PH, Halleen F. 2004. DNA phylogeny, morphology and pathogenicity of Botryosphaeria species on grapevines. *Mycologia* 96: 781–798.

- Pavlic D, Wingfield MJ, Barber P, Slippers B, Hardy GESTJ, Burgess TI. 2008. Seven new species of the Botryosphaeriaceae from baobab and other native trees in Western Australia. *Mycologia* 100: 851–866.
- Pérez CA, Wingfield MJ, Slippers B, Altier NA, Blanchette RA. 2010. Endophytic and canker-associated Botryosphaeriaceae occurring on non-native Eucalyptus and native Myrtaceae trees in Uruguay. *Fungal Diversity* 41: 53–69.
- Phillips AJL, Alves A, Abdollahzadeh J, Slippers B, Wingfield MJ, et al. 2013. The Botryosphaeriaceae: genera and species known from culture. *Studies in Mycology* 76: 51–167.
- Phillips AJL, Alves A, Correia A, Luque J. 2005. Two new species of Botryosphaeria with brown, 1-septate ascospores and Dothiorella anamorphs. *Mycologia* 97: 513–529.
- Phillips AJL, Alves A, Pennycook SR, Johnston PR, Ramaley A, et al. 2008. Resolving the phylogenetic and taxonomic status of dark-spored teleomorph genera in the Botryosphaeriaceae. *Persoonia* 21: 29–55.
- Rannala B, Yang Z. 1996. Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *Journal of Molecular Evolution* 43: 304–311.
- Rayner RW. 1970. A mycological colour chart. Kew, Surrey, UK: CMI and British Mycological Society.
- Rodríguez F, Oliver JF, Marin A, Medina JR. 1990. The general stochastic model of nucleotide substitutions. *Journal of Theoretical Biology* 142: 485–501.
- Slippers B, Boissin E, Phillips AJL, Groenewald JZ, Wingfield MJ, et al. 2013. Phylogenetic lineages in the Botryosphaeriales: A systematic and evolutionary framework. *Studies in Mycology* 76: 31–49.
- Slippers B, Crous PW, Denman S, Coutinho TA, Wingfield BD, Wingfield MJ. 2004. Combined multiple gene genealogies and phenotypic characters differentiate several species previously identified as Botryosphaeria dothidea. *Mycologia* 96: 83–101.
- Sutton BC. 1977. Coelomycetes VI. Nomenclature of generic names proposed for Coelomycetes. *Mycological Papers* 141: 1–253.
- Sutton BC. 1980. The Coelomycetes, Fungi imperfecti with acervuli, pycnidia and stromata. Commonwealth Mycological Institute, Kew, UK.
- Swofford DL. 2003. PAUP\* 4.0b10: Phylogenetic Analysis Using Parsimony (\*and other methods). Sinauer Associates, Sunderland, Massachusetts.
- Taylor K, Barber PA, Hardy GESTJ, Burgess TI. 2009. Botryosphaeriaceae from tuart (Eucalyptus gomphocephala) woodland, including descriptions of four new species. *Mycological Research* 113: 337–353.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. 1997. The ClustalX windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research* 25: 4876–4882.
- Urbez-Torres JR, Peduto F, Striegler RK, Urrea-Romero KE, Rupe JC, et al. 2012. Characterization of fungal pathogens associated with grapevine trunk diseases in Arkansas and Missouri. *Fungal Diversity* 52: 169–189.
- Wet J de, Slippers B, Preisig O, Wingfield BD, Tsopelas P, Wingfield MJ. 2009. Molecular and morphological characterization of Dothiorella casuarini sp. nov. and other Botryosphaeriaceae with diplodia-like conidia. *Mycologia* 101: 503–511.
- Young ND, Healy J. 2003. GapCoder automates the use of indel characters in phylogenetic analysis. *BMC Bioinformatics* 4: 6.
- Zhou S, Stanosz GR. 2001. Relationships among Botryosphaeria species and associated anamorphic fungi inferred from the analyses of ITS and 5.8S rDNA sequences. *Mycologia* 93: 516–527.